

# Chickenpox

Fact Sheet

## What is Chickenpox?

Chickenpox (varicella) is caused by varicella zoster virus (VZV). It is often a mild illness, but may be severe in infants, pregnant women, adults and persons with weakened immune systems. The greatest number of chickenpox cases occurs in the late winter and spring.

Before the chickenpox vaccine, almost everyone got chickenpox by adulthood--more than 95% of American adults have already had chickenpox.

## What are the symptoms of Chickenpox?

Chickenpox has a characteristic itchy rash that often starts on the trunk and face with pink spots and tiny fluid-filled blisters ("pox") that then dry and become scabs in 4 to 5 days.

The rash may be the first sign of illness and sometimes occurs with fever and general fatigue.

An infected person may have anywhere from only a few pox to more than 500 (average 300-400) pox on their body during the illness.

Chickenpox can sometimes have severe complications such as bacterial skin infections and pneumonia.

## How is Chickenpox spread?

Chickenpox virus is highly contagious and is spread by direct contact with secretions (such as saliva) or pox from an infected individual, or by respiratory secretions released into the air from sneezing or coughing. The illness starts 10 to 21 days after contact with an infected person.

## How is it spread (cont.)?

Infected persons are contagious from 1 to 2 days before the rash appears, until all the pox have formed scabs; usually 5 days from the time the pox appeared.

People with immune systems weakened by illness or drugs may require more time for their pox to scab over.

## How is Chickenpox diagnosed?

Chickenpox is typically diagnosed by the symptoms and the characteristic appearance of the rash.

Occasionally a healthcare provider will use laboratory tests for chickenpox if the diagnosis is unclear or if the illness is severe.

## How can Chickenpox be prevented?

Vaccination is the best way to prevent chickenpox.

- Children should receive two doses of varicella vaccine; the first at age 12-15 months and the second at age 4-6 years.
- Children 13 years and older and adults who do not have evidence of prior immunity routinely receive 2 doses of vaccine 4 to 8 weeks apart.
- Second dose catch-up vaccination is recommended for children, adolescents, and adults who previously received only 1 dose.
- The vaccine occasionally causes mild side effects including fever and a mild rash that can occur 5 to 26 days after immunization.
- You don't need the vaccine if you have already had chickenpox. If you are not sure whether you have had chickenpox, talk to your healthcare provider about getting a blood test.



Skagit County Public Health  
700 S 2nd St  
Mount Vernon, WA 98273

(360) 416-1500

Revised December 2018

Varicella vaccine is highly effective at preventing chickenpox, especially severe cases of the disease, and its complications.

### **What should I do if I've been exposed to chickenpox?**

If you are already immune to chickenpox, you most likely won't become sick. Most people do not get chickenpox more than once.

If you're not already immune to chickenpox, varicella vaccine given within 3 days of exposure can possibly prevent chickenpox. Vaccination 3 to 5 days after exposure probably won't prevent chickenpox but it may make it milder.

Persons at high risk for complications from chickenpox, such as immunocompromised persons, pregnant women, and premature infants, should contact their healthcare provider immediately.

### **What is "shingles"?**

Shingles, also called herpes zoster or zoster, is a painful skin rash caused by the varicella zoster virus (VZV). After a person recovers from chickenpox, the virus stays in the body. Usually the virus does not cause any problems; however, the virus can reappear years later, causing shingles.

Shingles usually starts as a rash on one side of the face or body. The rash starts as blisters that scab after 3 to 5 days. The rash usually clears within 2 to 4 weeks. Before the rash develops, there is often pain, itching, or tingling in the area where the rash will develop. Other symptoms can include fever, headache, chills and upset stomach.

Direct contact with the blisters of a person with shingles can cause chickenpox in someone who has never had chickenpox and has not received the varicella vaccine.

A single dose of shingles vaccine is indicated for adults 60 years of age and older. It can reduce the pain associated with shingles.

**For more information about chickenpox, please contact your health care provider or Skagit County Public Health at (360) 416-1500.**

### **More information available at:**

<https://www.cdc.gov/vaccines/parents/diseases/child/varicella.html>